

Suggested solutions for self-study & additional practice for the 3. Tutorial

Sample solution for the additional exercise 28:

- a) We use the function $\varphi(x) = k\sqrt{0.75k^2 + 2k + 1 + x}$. For the exact value we solve

$$s = k\sqrt{0.75k^2 + 2k + 1 + s} \quad \Rightarrow \quad s_1 = \frac{k}{2}(3k + 2), \quad s_2 = -\frac{k}{2}(k + 2).$$

The value s_2 is not of interest any more, thus $s = \frac{k}{2}(3k + 2)$.

For the analysis we get

$$\begin{aligned} \varphi'(x) &= \frac{k}{2\sqrt{0.75k^2 + 2k + 1 + x}} = \frac{k}{\sqrt{3k^2 + 8k + 4 + 4x}} \\ \varphi'(x_1) &= \frac{k}{\sqrt{9k^2 + 12k + 4}} \in [0, 1) \quad \forall k \in \mathbb{N} \end{aligned}$$

and we select e. g. the interval $I = [\alpha, \beta]$ with $-\frac{k^2}{2} - 2k - 1 < \alpha < \frac{k}{2}(3k + 2)$ and $\beta > \frac{k}{2}(3k + 2)$. Thus $|\varphi'(x)| < 1 \quad \forall x \in I$.

- b) We have to analyze the iteration $x^{(k+1)} = \varphi(x^{(k)})$ with $\varphi(x) = \cos x$.

The interval $I = [0, 1]$ is closed and it holds $\varphi(I) \subseteq I$ as \cos is monotonously falling on I and $\cos(0) = 1 \in I$ as well as $\cos(1) < 0.6 \in I$. Further holds $\max_{x \in I} |-\sin(x)| < 0.85 = L$. So the conditions of Banach's fixed-point theorem are satisfied and there exists a unique fixed-point in I and the iteration converges for all starting values $x_0 \in I$.

Sample solution for the additional exercise 29:

- a) Isolating x on the left hand side, we get

$$\ln(16 - x) = \sqrt{\frac{2}{3}x^2 + 4} \quad \Leftrightarrow \quad x = 16 - \exp\left(\sqrt{\frac{2}{3}x^2 + 4}\right),$$

hence the iteration is given by

$$\varphi_1(x) = 16 - \exp\left(\sqrt{\frac{2}{3}x^2 + 4}\right).$$

- b) Isolating x on the right hand side, we get

$$\ln(16 - x) = \sqrt{\frac{2}{3}x^2 + 4} \quad \Leftrightarrow \quad x = \pm \sqrt{\frac{3}{2}\left((\ln(16 - x))^2 - 4\right)}.$$

Since we are looking for an intersection in the interval $[1, 7]$, it is sufficient to choose the

iteration

$$\varphi_2(x) = \sqrt{\frac{3}{2} \left((\ln(16-x))^2 - 4 \right)}.$$

c) We compute the Lipschitz constants for φ_1 and φ_2 :

$$|\varphi_1'(x)| = \underbrace{\left| \exp \left(\sqrt{\frac{2}{3}x^2 + 4} \right) \right|}_{\geq \exp \left(\sqrt{\frac{2}{3}1^2 + 4} \right)} \cdot \underbrace{\left| \left(\frac{2}{3}x^2 + 4 \right)^{-\frac{1}{2}} \right|}_{\geq \left(\frac{2}{3}1^2 + 4 \right)^{-\frac{1}{2}}} \cdot \underbrace{\left| \frac{4}{3}x \right|}_{\geq \frac{4}{3}} \geq 1.9097 > 1,$$

$$|\varphi_2'(x)| = \underbrace{\left| 3 \ln(16-x) \right|}_{\leq 3 \ln(16-1)} \cdot \underbrace{\left| \frac{1}{16-x} \right|}_{(16-7)^{-1}} \cdot \underbrace{\left| \left(\frac{3}{2} \left((\ln(16-x))^2 - 4 \right) \right)^{-\frac{1}{2}} \right|}_{\leq \left(\frac{3}{2} \left((\ln(16-7))^2 - 4 \right) \right)^{-\frac{1}{2}}} \leq 0.81008 < 1.$$

Since $[1, 7]$ is a closed subspace of \mathbb{R} and $\varphi([1, 7]) \subset [1, 7]$, we are allowed to apply Banach's fixed-point theorem. Hence the second fixed-point iteration converges to $x = 2.095856$, whereas the first one does not.

d) For the iteration φ_2 a possible Lipschitz constant L is given by $L = 0.81008$. The a-priori estimate is

$$|x_i - x^*| \leq \frac{L^i}{1-L} |x_1 - x_0|,$$

hence we get the inequality

$$10^{-5} \leq \frac{0.81008^i}{0.18992} 1.033163.$$

We need at least 63 iterations to drop the distance $|x_i - x^*|$ below 10^{-5} .

Sample solution for the additional exercise 30:

a) We get the iteration

$$x_{k+1} = x_k - \frac{x_k^3 - 2x_k + 2}{3x_k^2 - 2}$$

and we obtain the following results

$$x_0 = -1, \quad x_1 = -4, \quad x_2 = -2.8261, \quad x_3 = -2.1467.$$

b) Starting at $x_0 = 1$, the first three iterations are given by

$$x_0 = 1, \quad x_1 = 0, \quad x_2 = 1, \quad x_3 = 0.$$

Therefore the sequence is periodical and does not converge to the zero of f .