

6. Additional tasks for exercise on „Introduction to Numerical Mathematics“

Problem 24:

Apply the power method as well as the inverse power method to

$$A = \begin{pmatrix} -2 & 1 & 0 \\ 1 & -3 & 1 \\ 0 & 1 & -4 \end{pmatrix}.$$

Calculate three steps starting from $x^{(0)} = 3^{-0.5}(1, 1, 1)^T$ for the power method as well as one step starting from $z^{(0)} = (1, -1, 0)^T$ for the inverse vector iteration. Give the approximations for the largest and smallest eigenvalues of A as well as the eigenvectors. Compare the approximations to the values

$$\lambda_1 = -4.7321, \quad \lambda_2 = -3.0, \quad \lambda_3 = -1.2679.$$

Problem 25:

Apply 3 steps of the shifted inverse power method for $\mu = -2.5$ to A from task 24. Use the starting vector $x^{(0)} = (1, 1, 1)^T$ and an LU -decomposition with pivoting of $A - \mu I$ with

$$L = \begin{pmatrix} 0.5 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0.8 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & -0.5 & 1 \\ 0 & 1.25 & -0.5 \\ 0 & 0 & -1.1 \end{pmatrix}.$$

Problem 26:

Consider once again the matrix A from task 24. Use a computer and

- (a) apply 7 steps of the QR-algorithm,
- (b) apply 7 steps of Jacobi's iteration.

Problem 27:

- (a) Apply the QR-algorithm to find all eigenvalues of the matrix $E = (E_{ij}) \in \mathbb{R}^{6 \times 6}$ defined by

$$E_{ij} = \begin{cases} 8 - i & \text{if } i = j, \\ -1 & \text{if } |i - j| = 1, \\ 0 & \text{otherwise.} \end{cases}, \quad i, j = 1, \dots, 6.$$

Determine the eigenvectors as well!

- (b) Apply Jacobi's method to compute the eigenvalues and eigenvectors of the matrix

$$A = (n + 1)^2 \text{tridiag}(1, -2, 1) \in \mathbb{R}^{n \times n}$$

for $n = 30$. Plot the eigenvectors associated to λ_i , $i \in \{1, 2, 29, 30\}$, against

$$x \in \mathbb{R}^n, \quad x_i = \frac{i}{n + 1}.$$