

7. Additional tasks for exercise on „Introduction to Numerical Mathematics“

Problem 28:

- (a) For  $k \in \mathbb{N}$  compute the value

$$s = \lim_{n \rightarrow \infty} k \sqrt[n]{\underbrace{\frac{3}{4}k^2 + 2k + 1 + k \sqrt{\frac{3}{4}k^2 + 2k + 1 + \dots + k \sqrt{\frac{3}{4}k^2 + 2k + 1}}}_{n \text{ times root}}}.$$

Specify a step function  $\varphi$  and an interval  $I$  such that the iteration  $x_{n+1} = \varphi(x_n)$  converges to  $s$  for all initial values  $x_0 \in I$ . Compute 5 iterations with a proper starting value.

What is the exact value of  $s$  for a fixed  $k$ ?

- (b) If you enter a number  $x \in (0, \pi)$  into a calculator and press the "cos" key several times, you observe numerically a convergence to a fixed-point. Analyze this behavior with Banach's fixed-point theorem (Specify an interval, check preconditions, specify Lipschitz constant).

Problem 29:

Compute the intersection of the curves  $f(x) = \ln(16 - x)$  and  $g(x) = \sqrt{\frac{2}{3}x^2 + 4}$  in  $[1, 7]$ . To this end use a fixed-point iteration

- (a) by isolating  $x$  on the left hand side of  $f(x) = g(x)$ ,  
(b) by isolating  $x$  on the right hand side of  $f(x) = g(x)$ .  
(c) Analyze both iterations in terms of convergency.  
(d) How many iterations are necessary at the latest to drop with the distance  $|x_i - x^*|$  below a level of  $\varepsilon = 10^{-5}$ ?

Problem 30:

Apply Newton's method to solve  $f(x) = x^3 - 2x + 2 = 0$ . Perform three steps for each start value.

- (a) Use  $x_0 = -1$ .  
(b) Use  $x_0 = 1$ .